



Nilocin Wart & Verruca Treatment Pen

UK



For external use only
Keep out of reach of children
Nilocin® contains a corrosive agent.



Read this instruction carefully prior to use

INDICATION

Nilocin® Wart Pen is indicated for the treatment of warts and verrucas found on the feet, hands, body arms and legs. The product can be used by adults and children from 4 years old.



WARTS AND VERRUCA

Warts and verrucas (hereafter defined as warts) are local growths in the skin caused by the human papilloma virus (HPV). They may be found alone or in clusters. There exist different types of warts, most commonly found on hands, feet, elbows and knees. Warts are contagious and spread by contact of the virus with damaged skin.

NILOCIN®

The Nilocin Wart treatment is a CE marked Medical Device. The pack comprises a treatment pen and this instruction leaflet.

The Nilocin Wart Pen contains 3 ml, sufficient for minimum 200 applications. Nilocin® is specifically designed for topical administration directly on warts.

Nilocin® is immediately absorbed into the wart. The pen contains a keratolytic compound which breaks down tissue locally and thereby eliminates warts. Nilocin® can be used for treatment of children as young as 4 years of age.



WARNINGS & PRECAUTIONS

Only for external use. Keep out of reach of children. The product contains a corrosive agent with a strong smell. Do not inhale.

- Do NOT use the product for genital warts or in the facial area.
- Do not use Nilocin® if you are known to be hypersensitive or allergic to the ingredients, as well as bee/ant stings.
- When applying, avoid contact with healthy skin surrounding the wart.
- Do not expose the treated area to sunlight, until the skin has returned to normal.
- Do not apply Nilocin® to sore or bleeding warts, irritated, inflamed, blistered or damaged skin. Pause until the skin has recovered.
- Do not use on warts with hairs, birthmarks, dark moles, pigmented skin or in case you have doubts if the skin condition is a wart.
- Extra care should be taken when applying the treatment to thin and sensitive skin, and when treating children and people with diabetes.

It is better to apply too little liquid if in doubt. Over application can result in side effects such as wounds, blisters, inflammation, tissue death (necrosis), scars, and chemical burns

TECHNICAL DATA - Product name: nilocin Product: Wart & Verruca Treatment Pen **Ingredients:** Formic acid, Dimethyl isosorbide, Aqua, Caprylic acid, Oenonium Vulgare leaf Extract. **Content:** 3 ml, sufficient for minimum 200 applications. **Registration:** Nilocin® is classified and registered as a Medical Device Class IIa. **Storage:** Store at 2-25 degree Celsius, in the original packaging, out of direct sunlight and out of reach and sight of children. Not to be used after the expiry date. **Manufacturer:** Unigroup AP5, Diplomvej 373, DK-2800 Lyngby, Denmark. www.nilocin.com. Information updated: 2019-07

- Do not exceed the number of recommended applications.
- Consult your doctor if the treated area becomes red and painful or gets inflamed.
- Single patient - multiple use
- Do not use in combination with other wart removal therapies.
- Do not use for children under 4 years of age, unless directed by doctor.

SHOULD THE PRODUCT ACCIDENTALLY COME INTO CONTACT WITH THE EYES OR MUCOSAL MEMBRANES, RINSE WITH PLENTY OF WATER FOR 15 MINUTES AND SEEK MEDICAL ADVICE

HOW TO USE NILOCIN

It is recommended to soak the wart in luke warm water for at least 5 minutes to soften the skin prior to treatment. If the wart is covered by hard skin, gently scrape the surface of the wart (e.g. with a scissors blade) and carefully remove the hard skin. Do not file the wart. It is recommended that adults apply the treatment to children.

- Remove the cap from the Nilocin pen by pressing and turning the cap counterclockwise
- Gently massage the wart with the tip of the treatment pen for a maximum of 20 seconds. The solution penetrates immediately. Make sure to cover the whole wart and take care not to touch or spill onto surrounding skin. Recap the pen firmly after application.
- Repeat the treatment 3 times a week, a day apart, for up to 6 weeks. **Do not exceed the number of recommended weekly applications.**

The Nilocin pen delivers a small amount of a colorless solution. If in doubt whether the tip of the pen is dry, press the tip against a piece of absorbent paper to check the liquid flow.

Should any discomfort or pain occur, stop application. If it burns, sting, or if serum is applied to healthy skin, rinse with cold water.

POSSIBLE SIDE EFFECTS

It is important to use the product as instructed. Do not overexpose as this can cause skin damage (please see **PRECAUTIONS** above).

A mild tingling or burning sensation can in some cases be felt for 10-15 minutes. If overexposed temporary side effects can occur, such as stinging, redness, pain or bleeding. Pause treatment until the skin has fully recovered. There are no known risks for pregnant or breastfeeding women.

STORAGE

Store between 2 to 25 degree Celsius and out of direct sunlight, enclosed in the original packaging and out of the reach of children. Do not use after the expiry date, or 6 months after removing the cap from the pen for the first time. Carefully recap the pen after use.

The product can be disposed with the household rubbish.

Unigroup AP5, Diplomvej 373, DK-2800 Lyngby, Denmark
Nilocin is a registered trademark of Unigroup AP5.



UG201702-101034-UK



Nilocin Wart & Verruca Treatment Pen

UK



For external use only
Keep out of reach of children
Nilocin® contains a corrosive agent.



Read this instruction carefully prior to use

INDICATION

Nilocin® Wart Pen is indicated for the treatment of warts and verrucas found on the feet, hands, body arms and legs. The product can be used by adults and children from 4 years old.



WARTS AND VERRUCA

Warts and verrucas (hereafter defined as warts) are local growths in the skin caused by the human papilloma virus (HPV). They may be found alone or in clusters. There exist different types of warts, most commonly found on hands, feet, elbows and knees. Warts are contagious and spread by contact of the virus with damaged skin.

NILOCIN®

The Nilocin Wart treatment is a CE marked Medical Device. The pack comprises a treatment pen and this instruction leaflet.

The Nilocin Wart Pen contains 3 ml, sufficient for minimum 200 applications. Nilocin® is specifically designed for topical administration directly on warts.

Nilocin® is immediately absorbed into the wart. The pen contains a keratolytic compound which breaks down tissue locally and thereby eliminates warts. Nilocin® can be used for treatment of children as young as 4 years of age.



WARNINGS & PRECAUTIONS

Only for external use. Keep out of reach of children. The product contains a corrosive agent with a strong smell. Do not inhale.

- Do NOT use the product for genital warts or in the facial area.
- Do not use Nilocin® if you are known to be hypersensitive or allergic to the ingredients, as well as bee/ant stings.
- When applying, avoid contact with healthy skin surrounding the wart.
- Do not expose the treated area to sunlight, until the skin has returned to normal.
- Do not apply Nilocin® to sore or bleeding warts, irritated, inflamed, blistered or damaged skin. Pause until the skin has recovered.
- Do not use on warts with hairs, birthmarks, dark moles, pigmented skin or in case you have doubts if the skin condition is a wart.
- Extra care should be taken when applying the treatment to thin and sensitive skin, and when treating children and people with diabetes.

It is better to apply too little liquid if in doubt. Over application can result in side effects such as wounds, blisters, inflammation, tissue death (necrosis), scars, and chemical burns

TECHNICAL DATA - Product name: nilocin Product: Wart & Verruca Treatment Pen **Ingredients:** Formic acid, Dimethyl isosorbide, Aqua, Caprylic acid, Oenonium Vulgare leaf Extract. **Content:** 3 ml, sufficient for minimum 200 applications. **Registration:** Nilocin® is classified and registered as a Medical Device Class IIa. **Storage:** Store at 2-25 degree Celsius, in the original packaging, out of direct sunlight and out of reach and sight of children. Not to be used after the expiry date. **Manufacturer:** Unigroup AP5, Diplomvej 373, DK-2800 Lyngby, Denmark. www.nilocin.com. Information updated: 2019-07

- Do not exceed the number of recommended applications.
- Consult your doctor if the treated area becomes red and painful or gets inflamed.
- Single patient - multiple use
- Do not use in combination with other wart removal therapies.
- Do not use for children under 4 years of age, unless directed by doctor.

SHOULD THE PRODUCT ACCIDENTALLY COME INTO CONTACT WITH THE EYES OR MUCOSAL MEMBRANES, RINSE WITH PLENTY OF WATER FOR 15 MINUTES AND SEEK MEDICAL ADVICE

HOW TO USE NILOCIN

It is recommended to soak the wart in luke warm water for at least 5 minutes to soften the skin prior to treatment. If the wart is covered by hard skin, gently scrape the surface of the wart (e.g. with a scissors blade) and carefully remove the hard skin. Do not file the wart. It is recommended that adults apply the treatment to children.

- Remove the cap from the Nilocin pen by pressing and turning the cap counterclockwise
- Gently massage the wart with the tip of the treatment pen for a maximum of 20 seconds. The solution penetrates immediately. Make sure to cover the whole wart and take care not to touch or spill onto surrounding skin. Recap the pen firmly after application.
- Repeat the treatment 3 times a week, a day apart, for up to 6 weeks. **Do not exceed the number of recommended weekly applications.**

The Nilocin pen delivers a small amount of a colorless solution. If in doubt whether the tip of the pen is dry, press the tip against a piece of absorbent paper to check the liquid flow.

Should any discomfort or pain occur, stop application. If it burns, sting, or if serum is applied to healthy skin, rinse with cold water.

POSSIBLE SIDE EFFECTS

It is important to use the product as instructed. Do not overexpose as this can cause skin damage (please see **PRECAUTIONS** above).

A mild tingling or burning sensation can in some cases be felt for 10-15 minutes. If overexposed temporary side effects can occur, such as stinging, redness, pain or bleeding. Pause treatment until the skin has fully recovered. There are no known risks for pregnant or breastfeeding women.

STORAGE

Store between 2 to 25 degree Celsius and out of direct sunlight, enclosed in the original packaging and out of the reach of children. Do not use after the expiry date, or 6 months after removing the cap from the pen for the first time. Carefully recap the pen after use.

The product can be disposed with the household rubbish.

Unigroup AP5, Diplomvej 373, DK-2800 Lyngby, Denmark
Nilocin is a registered trademark of Unigroup AP5.



UG201702-101034-UK